

## Relationship of weight and height with waist circumference, body mass index and conicity index in adolescents\*

Betty M Pérez<sup>1</sup>, Maritza Landaeta-Jiménez<sup>2</sup>

<sup>1</sup>Central University of Venezuela "Rodolfo Quintero", Economic and Social Sciences Institute, Caracas, Venezuela

<sup>2</sup>Fundacredesa, Department of Auxology, Division of Biological Research, Caracas, Venezuela

### Summary

Obesity or overweight in adolescents are a public health issue. Waist circumference (WC) body mass index (BMI), and conicity index (CI) have been used to assess these conditions and the risks involved. The aim of this study was to assess the influence of weight and height on WC, BMI and CI in adolescents. Student *t* test, ANOVA, Pearson correlations and binary correspondence were used. At 13-14 years, BMI ( $r = 0.89$ ;  $r = 0.92$ ) and WC ( $r = 0.85$ ;  $r = 0.83$ ) showed high correlations with weight. C had a low correlation with weight and lesser dependence on height. Binary correspondence analysis, showed that high values of weight were associated with high values of CI. In this sample, CI is not an indicator of body mass as a whole, yet it shows that heavy weight is associated with centripetal fat distribution as measured by this indicator and in this sense, it can be used as an index of fat distribution.

**Key words** waist circumference, body mass index, conicity index, body composition, adolescents, Venezuela

### Introduction

Obesity and/or overweight in children and adolescents are a matter of concern in public health, both at individual and population levels. Overweight in young age is associated with overweight in adulthood (1,2). This condition affects about 33% of USA adult population (3), and the prevalence in youths has increased in the latest national health survey (4). The same tendency is also found in Venezuela where by 1999, the condition has reached 13% of youths (5). Furthermore, the nutrition transition in low-income countries, a condition where under and overnutrition exist simultaneously, is occurring very rapidly (6).

Screening over time for overweight and pattern of fat distribution is becoming an important aspect of pediatric care, given its potential long term health implications (7-9). In recent years the height-weight ratio, expressed as body mass index (BMI;  $\text{kg}/\text{m}^2$ ) is increasingly used as a reliable and valid indicator of obesity (10,11). Nevertheless, studies conducted in adults have also advocated that waist circumference (WC) and conicity index (CI), are valid indicators of obesity and risk factors of cardiovascular disease (12-14).

The evidence for children and youths is less compelling for BMI, WC and CI. Flodmark et al. (15) support the idea that WC is somehow related to a potentially atherogenic lipoprotein profile in obese children, whereas Freedman et al. (7) found, in the Bogalusa Heart Study, that the prevalences for dyslipidemia, hyperinsulinemia and hypertension increased at the same time as

\*Presented at the IX International Congress of Auxology 3<sup>rd</sup>-6<sup>th</sup> September 2000, Turin, Italy

BMI. On the other hand, data on conicity are limited. However, CI is associated with risk the of high levels of tryglicerides in Venezuelan females adolescents (16) and with other risk factors of coronary and heart diseases (14,17).

Although the ideal index to assess leanness and obesity should strongly correlate with body weight and minimize the influence of height (18), there is no clear consensus on what method should be used in children and adolescents, considering individual differences in maturity, timing and tempo. Therefore, the aim of this study was to assess the mediating influence of weight and height on WC, BMI and CI in Venezuelan adolescents, 11-16 years of age, seeking the best index to evaluate body composition at these ages.

## Materials and methods

The data were obtained from a cross-sectional study of 394 adolescents, 11-16 years of age from Caracas, capital city of Venezuela. The sample included 186 males and 208 females. Environmental, cultural and social conditions of the families of the youths included in the study classified them as of low socioeconomic status. This integrated analysis concerning growth, nutritional status and body composition, along with socioeconomic factors in this sample, has been reported previously (19,20).

### Anthropometry

The survey was carried out by a well trained group with quality controls being provided during the field work. The anthropometrical measurements were made in the morning, using portable equipment and following standard procedures (21); individuals wore minimal clothing. Height was appreciated to the nearest 0.1 cm with the subject in standing position, bare footed and with the head held in the Frankfort plane. A gentle pressure was applied on the mastoid process during measurement. On the other hand, weight was measured to the nearest 0.1 kg.

Minimum waist, taken directly on the skin, was made to the nearest 0.1 cm, using a flexible non-elastic steel tape\* with the subject standing, at the minimal abdominal girth, approximately midway between the xiphoid process and the umbilicus.

BMI was assessed as weight (kg) divided by the square of height (m) (22). CI was calculated according to the formula of Valdez et al. (23), as follows:

$$C = \frac{WC (m)}{0.109\sqrt{wt (kg)/ht (m)}}$$

as a measure of central adiposity, is based on quantifying the deviation from the circumference of an imaginary cylindrical shape derived from the weight and height of the individual. In comparison with other indices, it has been claimed to be easier to obtain and requiring less undressing besides being independent of hip circumference (24).

### Statistical analysis

Statistical analysis was carried out with the SPSS program (25). The sample was divided into three age groups: 11-12, 13-14 and 15-16 years. Analytic methods included Student t tests to identify sexual differences, and one way analyses of variance (ANOVA), to test for differences between age groups within each sex. In addition, Scheffé post hoc test was used to identify which pairs of age groups differed significantly. Two linear regressions were also performed with height and weight as independent variables. The line of best fit was determined by least squares method. Analysis of correspondence (26) was also performed to measure the degree of association among categories defined according to references and constructed intervals. Weight, height and BMI were classified as low ( $x < P10$ ), average ( $P10 \geq x \leq P90$ ) and high ( $x > P90$ ) according to percentile cut off points of a national reference (27). After verifying normality with the Kolmogorov-Smirnov-Lilliefors test, ( $p > 0.05$ ), and in absence of national references, toler-

\* Hoechst mass, West Germany

**Table I** Descriptive anthropometric statistics by age and sex

Age groups	No.	Weight	Height	WC	BMI	CI
<i>Boys</i>						
<b>11-12</b>						
Mean	70	34.44*	141.26*	60.21	17.16	1.12*
CV (%)		20.05	5.36	9.81	14.51	4.78
<b>13-14</b>						
Mean	64	46.59	156.95	64.44	18.66*	1.10*
CV (%)		24.70	6.33	12.00	17.03	5.82
<b>15-16</b>						
Mean	52	54.94	165.57*	67.78	19.90*	1.09*
CV (%)		17.91	5.01	8.91	12.03	5.52
<i>Girls</i>						
<b>11-12</b>						
Mean	80	37.12*	145.21*	59.03	17.49	1.08*
CV (%)		19.34	5.20	8.76	13.74	4.71
<b>13-14</b>						
Mean	75	48.68	155.00	64.60	20.16*	1.06*
CV (%)		18.87	3.78	10.59	14.85	5.65
<b>15-16</b>						
Mean	53	52.55	155.44*	67.18	21.72*	1.06*
CV (%)		12.21	3.28	7.27	10.07	4.37
<i>Legend:</i> WC = waist circumference; BMI = body mass index; CI = conicity index CV (%): coefficient of variation in percentage * ( $p < 0.05$ ) sexual dimorphism						

ance intervals for WC and CI were constructed and classified, as low ( $< \bar{x} + s_x$ ), average ( $\bar{x} \pm s_x$ ) and high ( $> \bar{x} + s_x$ ).

## Results

Table I presents means and coefficients of variations for weight, height, WC, BMI and CI by age groups and sex. All variables values except CI increased with age. Also they differed significantly between sexes, specially CI that attained differences at all age groups. On the other hand WC, was the only variable that did not show sexual dimorphism.

Results of ANOVA and Scheffé tests (not shown in the text) indicated differences by age groups as follows:

a) weight and WC in all age groups in both sexes;

- b) BMI in boys and in girls at 11-12 years compared to the other two age groups;  
 c) CI differences in boys between the younger and the two older groups, but no differences in girls;  
 d) height differs significantly in boys in all age groups;  
 e) the two older age groups of girls do not differ in height, BMI and CI.

Tables II and III depict results of the linear regression, with weight and height as independent variables. WC, BMI and CI were first regressed onto weight, and the resulting estimated coefficient models were statistically significant for both, WC and BMI ( $p < 0.05$ ). In both sexes WC had a high significant correlation with weight, ( $r = 0.73$  to  $0.85$ ) and BMI ( $0.83$  to  $0.92$ ) in boys and girls. On the other hand, WC and BMI were related to height to a lesser extent, but the correlations were significant for WC ( $0.35$  to

**Table II** Results of simple linear regression of waist circumference, body mass index and conicity index on weight and height, by age group in boys

Age groups	Independent variable: Weight				Independent variable: Height				
	Estimate	p-value	r	r <sup>2</sup>	Estimate	p-value	r	r <sup>2</sup>	
<b>Waist circumference (WC)</b>									
11-12	b <sub>0</sub>	35.798	0.000	0.829*	0.687	21.432	0.092	0.352*	0.124
	b <sub>1</sub>	0.709	0.000			0.274	0.003		
13-14	b <sub>0</sub>	37.841	0.000	0.840*	0.722	4.696	0.730	0.489*	0.239
	b <sub>1</sub>	0.571	0.000			0.381	0.000		
15-16	b <sub>0</sub>	43.049	0.000	0.734*	0.538	23.146	0.150	0.371*	0.137
	b <sub>1</sub>	0.450	0.000			0.270	0.007		
<b>Body mass index (BMI)</b>									
11-12	b <sub>0</sub>	6.742	0.000	0.839*	0.703	8.613	0.125	0.184	0.034
	b <sub>1</sub>	0.302	0.000			0.06050	0.128		
13-14	b <sub>0</sub>	7.184	0.000	0.892*	0.795	-3.919	0.495	0.450*	0.202
	b <sub>1</sub>	0.246	0.000			0.144	0.000		
15-16	b <sub>0</sub>	8.393	0.000	0.861*	0.741	1.481	0.813	0.386*	0.149
	b <sub>1</sub>	0.209	0.000			0.111	0.005		
<b>Conicity index (CI)</b>									
11-12	b <sub>0</sub>	1.113	0.000	0.043	0.002	1.259	0.000	0.134	0.018
	b <sub>1</sub>	0.00034	0.721			-0.00095	0.268		
13-14	b <sub>0</sub>	1.097	0.000	0.006	0.000	1.312	0.000	0.215	0.046
	b <sub>1</sub>	-0.00003	0.963			-0.00138	0.088		
15-16	b <sub>0</sub>	1.135	0.000	0.148	0.022	1.449	0.000	0.304*	0.092
	b <sub>1</sub>	-0.00090	0.293			-0.00219	0.028		

Legend:  $Y_1 = b_0 + b_1 \text{weight}$ ,  $Y_2 = b_0 + b_1 \text{height}$

\* Correlate statistically significant

0.49), but not consistent for BMI (0.01 to 0.45). CI had negligible correlations with weight (0.01 to 0.15), and slightly higher with height (0.07 to 0.30).

At 13-14 years, weight accounted for 72-69% of waist variability and for 80-85% of BMI variability, while regression of height was not good enough to explain variability on these variables. Weight and height, on the other hand, were not strong enough to explain the variability of the CI.

When the indices are plotted in relation to body weight and height, based on the best regression fit, the best r<sup>2</sup> are attained at 13-14 years of age in boys and girls, r<sup>2</sup> = 0.795 and 0.847, for BMI and WC respectively. On the other hand, in this group WC ap-

pears as the most dependable variable on height, with r<sup>2</sup> of 0.239 and 0.141, for boys and girls. Generally speaking, boys and girls were similar in terms of patterns followed by the regression lines of best fit, describing the plot of the variables for the age-groups against weight (tables II, III and figure 1).

These results suggest that in the sample of 13-14 years old adolescents, BMI met the criteria as a satisfactory index, in that there was a strong correlation with weight, and a negligible correlation with height. The second best index according with this criteria was WC. CI does not show a satisfactory relationship owing to the structure of the index itself.

**Table III** Results of simple linear regression of waist circumference, body mass index and conicity index on weight and height, by age group in girls

Age groups	Independent variable: Weight				Independent variable: Height				
	Estimate	p-value	r	r <sup>2</sup>	Estimate	p-value	r	r <sup>2</sup>	
<b>Waist circumference (WC)</b>									
11-12	b <sub>0</sub>	37.055	0.000	0.822*	0.676	15.989	0.120	0.433*	0.187
	b <sub>1</sub>	0.592	0.000			0.296	0.000		
13-14	b <sub>0</sub>	34.582	0.000	0.828*	0.685	-0.378	0.864	0.376*	0.141
	b <sub>1</sub>	0.617	0.000			0.439	0.001		
15-16	b <sub>0</sub>	35.220	0.000	0.799*	0.639	15.835	0.421	0.346*	0.119
	b <sub>1</sub>	0.608	0.000			0.330	0.011		
<b>Body mass index (BMI)</b>									
11-12	b <sub>0</sub>	6.182	0.000	0.859*	0.738	4.378	0.387	0.284*	0.080
	b <sub>1</sub>	0.288	0.000			0.09030	0.011		
13-14	b <sub>0</sub>	5.558	0.000	0.920*	0.847	-4.815	0.586	0.315*	0.100
	b <sub>1</sub>	0.300	0.000			0.161	0.006		
15-16	b <sub>0</sub>	6.821	0.000	0.832*	0.692	20.914	2.243	0.012	0.000
	b <sub>1</sub>	0.284	0.000			0.00521	0.087		
<b>Conicity index (CI)</b>									
11-12	b <sub>0</sub>	1.118	0.000	0.153	0.023	1.245	0.000	0.172	0.030
	b <sub>1</sub>	-0.00108	0.175			-0.00116	0.127		
13-14	b <sub>0</sub>	1.033	0.000	0.091	0.008	1.170	0.000	0.068	0.005
	b <sub>1</sub>	0.00059	0.440			-0.00069	0.563		
15-16	b <sub>0</sub>	1.004	0.000	0.153	0.023	0.827	0.000	0.166	0.028
	b <sub>1</sub>	0.00101	0.274			0.00151	0.235		

Legend:  $Y_1 = b_0 + b_1 \text{weight}$ ,  $Y_2 = b_0 + b_1 \text{height}$


\* Correlate statistically significant

The factorial axis of the correspondence analysis (figures 2, 3 and 4) showed the pattern of behavior of WC, BMI and CI adjusted to weight and height. Figure 2 showed the typical U-shaped associations of waist circumference that were similar and consistent, but of variable statistical significance for weight in boys and girls, respectively, in all age groups. For height there was a near direct association for the total sample in both sexes. However, in girls, when the information is analyzed by age, this pattern is discontinued for low values.

BMI (figure 3) was also in close association with weight in both sexes as a whole (U-shape) and by age groups. Similarly, although less strictly, BMI was related to

height in boys, especially in the 13-14 year age group. In girls the BMI acted more independently on height, although high values of height are found in correspondence with high values of the BMI.

The CI (figure 4) showed low association with weight for the whole group in both boys and girls. However, in 13-14 year old girls, there was a correspondence between high values of CI and weight. On the other hand referring to height, there was an inverse relationship in boys; it was found that high values of CI were associated with low height values and viceversa. That is to say that the boys of our sample, aged 13-14 years and with low height tend to be more centripetal in their body fat distribution.



**Figure 1** Scatterplots of waist circumference, body mass and conicity index, adjusted by height and weight in 13-14 year old Venezuelan youths.

## Discussion

Overweight and/or obesity in children and adolescents are a prevalent worldwide multifactorial health problem, whose causes have been attributed to different etiologies. Gene-environment-culture interaction appears important. A list of environmental causes has been outlined: eating disorders, away-from-home eating, dietary energy consumption that overcomes energy expenditure, lack of energy expenditure through physical activities, among others (3,28). The present results show that WC is closely

related to weight ( $r^2 = 0.722$ ;  $r^2 = 0.685$ ) for boys and girls specifically at 13-14 years of age. WC showed a similar pattern for both sexes, having strong correlation with weight and moderate although statistically significant correlation with height.

Studies carried out in adult males and females in The Netherlands suggest that waist circumference is gaining support as a good anthropometric indicator to predict fatness and is of clinical and public health use, (13). In addition it is a simple yet sensitive method for screening both overweight and topography of fat distribution, when accurately classifying 83% of the female sample aged 16-80 years (29). A review made by Vanitallie (30) over a period from 1914 up to date, reports the reliability of WC, both as a measure of adiposity and a predictor of risk of obesity related to metabolic and cardiovascular diseases.

Several studies have examined at the correlations of BMI with weight, and Roche et al. (31) found that this relationship is a good predictor of obesity, based on estimated body fat. The long-term follow-up studies of BMI, like that of Rolland-Cachera et al. (32), indicated that BMI accounts more than skinfold measurements. Cole (22), also noted that BMI is a very useful tool to identify both obesity and malnutrition in children and adults. However, owing to the very skew distribution of the index during childhood and early adulthood, (even skewer than weight); the index should be transformed in order to approximate to a more normal distribution.

A primary finding in the current analysis confirms that BMI scaled strongly ( $r^2 = 0.847$ ,  $p < 0.05$ ) and positively ( $b = 0.30$ ) to weight (table III, figure 1). Although BMI includes a mixture of both fat mass and lean body mass (18,33) the positive slope indicate that a higher BMI is attained at increasing levels of weight. In boys more pronounced U-shaped curves were apparent for both WC and BMI in relation to weight. The results reported by Pietrobelli et al. (10), who worked with an unadjusted association of






Figure 2 Factorial axis of the correspondence analysis for waist circumference referred to weight and height. Total groups and selected ages.



**Figure 3** Factorial axis of the correspondence analysis for body mass index referred to weight and height. Total groups and selected ages.





**Figure 4** Factorial axis of the correspondence analysis for body conicity index referred to weight and height. Total groups and selected ages.

BMI, with total body fat and percentage of body fat, also pointed out a strong association between BMI and total body fat.

C showed a poor relation to weight ( $r^2 = 0.008$ ;  $b = 0.005$ ) in this sample of Venezuelan adolescents, yet, at the same time, this fact also suggests a negligible dependence of height ( $r^2 = 0.018$ ;  $b = -0.00095$ ), although in this group, boys with low height showed some tendency towards centripetal distribution, and high height values were associated with peripheral shape. In this group of Venezuelan adolescents, CI is not a good predictor of overweight. However, a closer insight through correspondence analysis revealed, particularly in girls, some close correlation between abdominal adiposity, as measured by the CI, and high values of weight. These findings could lead to investigate maturation related effects in their body composition.

### Acknowledgements

This study was given financial assistance by Consejo de Desarrollo Científico y Humanístico from the Central University of Venezuela. We gratefully acknowledge the statistical job performed by Miss Ynay Escalante. Also we appreciate the revision of the English manuscript made by Dr. Robert Malina.

**Correspondence** Betty M Pérez, Apartado de Correos 78162, La Urbina, Caracas 1074, Venezuela; e-mail: mariusa@telcel.net.ve

### References

- Guo SS, Roche AF, Chulmea WC, Gardner JC, Siervogel RM. The predictive value of childhood body mass index values for overweight at age 35. *Am J Clin Nutr* 1994;59:810-19.
- Rossner S. Childhood obesity and adulthood consequences. *Acta Paediatr* 1998;87:1-5.
- Schonfeld-Warden N, Warden CH. Obesidad en niños: Consideraciones sobre sus causas y tratamientos. *Clín Ped Norteam*. Vol 2. Endocrinología. Mc Graw-Hill Interamericana, 1997.
- Troiano R, Flegal K.M, Kuczmarski RJ, Campbell SM, Johnson CL. Overweight prevalence and trends for children and adolescents. *Arch Pediatr Adolesc Med* 1995;149:1085-91.
- Instituto Nacional de Nutrición. Valores de referencia de energía y nutrientes para la población venezolana. Cuaderno Azul 2001;53.
- Popkin BM. The nutrition transition in low-income countries: an emerging crisis. *Nutr Rev* 1994;52:285-98.
- Freedman D, Dietz, W D, Srinivasan SR, Berenson G. The relation of overweight to cardiovascular risk factors among children and adolescents: The Bogalusa Heart Study. *Pediatrics* 1999;103:1175-82.
- Dietz WH. Childhood weight affects adult morbidity and mortality. *J Nutr* 1998;128:411-4.
- Caprio S. Relationship between abdominal visceral fat and metabolic risk factors in obese adolescents. *Am J Hum Biol* 1999;11:259-66.
- Pietrobelli A, Faith M, Allison D, Gallagher D, Chiumello G, Heymsfield S. Body mass index as a measure of adiposity among children and adolescents: A validation study. *J Pediatr* 1998;132:204-10.
- Daniels SR, Khoury PR, Morrison JA. The utility of body mass index as a measure of body fatness in children and adolescents: differences by race and gender. *Pediatrics* 1997;99:804-7.
- Seidell JC, Cigolini M, Charzewska J, Ellsinger BM, Deslypere JP, Cruz A. Fat distribution in European men: a comparison of anthropometric measurements in relation to cardiovascular risk factors. *Int J Obes* 1992;16:17-22.
- Lean MEJ, Han TS, Morrison CE. Waist circumference as a measure for indicating need for weight management. *BMJ* 1995;311:158-61.
- Mueller WH, Meinger JC, Liehr P, Chan W, Chandler PS. Conicity: a new index of body fat distribution-what does it tell us. *Am J Hum Biol* 1996;8:489-96.
- Flodmark CE, Sveger T, Nilsson-Ehle P. Waist measurement correlates to a potentially atherogenic lipoprotein profile in obese 12-14-year-old children. *Acta Paediatr* 1994;83:941-5.
- Pérez B, Landaeta-Jiménez M, Ledezma T, Mancera A. Adiposidad, distribución de grasa y lípidos séricos en adolescentes venezolanos. In: Tito A. Varela, ed. *Investigaciones en Biodiversidad Humana*. Santiago de Compostela: Publicaciones de la Universidad de Santiago de Compostela 2000;539-46.
- Valdez R, Seidell JC, Ahn YI, Weiss KM. A new index of abdominal adiposity as an indicator of risk for cardiovascular disease. A cross-population study. *Int J Obes* 1992;16:77-82.
- Gasser Th, Kneip A, Ziegeer P, Molinari L, Prader A, Largo RH. Development and outcome of indices of obesity in normal children. *Ann Hum Biol* 1994;21:275-86.
- Ledezma T, Pérez B, Landaeta-Jiménez M. Pobreza y desnutrición: Factores limitantes del desarrollo humano. *Contribuciones a la Antropología Física Latinoamericana* 1995;1:19-29.
- Pérez B, Landaeta-Jiménez M, Ledezma T. Social and biological profiles of children at risk in a

- Venezuelan community. In: Human biology: Global Developments. Patiala: India, 1996:197-216.
21. Weiner JS, Lourie J A. Human biology: a guide to field methods In: Tanner JM, Hiernaux J, Jarman S. Growth and physique. Blackwell Scientific Publications: Oxford, 1969.
  22. Cole JE. Weight-Stature indices to measure underweight, overweight, and obesity. In: Anthropometric assessment of nutritional status. Wiley-Liss Inc, 1991:83-111.
  23. Valdez R. A simple model-based index of abdominal adiposity. *J Clin Epidemiol* 1991;44:955-6.
  24. Wardle J, Wrihston K, Gibson L. Body fat distribution in South Asian women and children. *Int J Obesity* 1996;20:267-71.
  25. Statistical Software System. SPSS 7.5. SPSS Inc, 1997.
  26. Greenacre M. Correspondence analysis in practice. Academic Press: London, 1993.
  27. López Contreras de Blanco M, Landaeta-Jiménez M, Espinoza I, Torralba C. Crecimiento físico y maduración. Vol 2. In: Estudio nacional de crecimiento y desarrollo de la república de Venezuela. FUNDA-CREDESA: Venezuela, 1996:407-619.
  28. Mumbiela PB, Sanmartin ZS, González AC. Obesidad infantil y hábitos alimentarios. *Rev Enferm* 1997;20:11-7.
  29. Taylor R, Keil D, Gold EJ, Williams S, Goulding A. Body mass index, waist girth, and waist-to-hip ratio as indexes of total and regional adiposity in women: evaluation using receiver operatin characteristic curves *Am J Clin Nutr* 1998;67:44-9.
  30. Vanitallie T. Waist circumference: a useful index in clinical care and health promotion. *Nutr Rev* 1998; 56:300-13.
  31. Roche AF, Siervogel RM, Chulmea WC, Webb P. Grading body fatness from limited anthropometric data. *Am J Clin Nutr* 1981;34:2831-7.
  32. Rolland-Cachera M, Bellisle F, Deheeger M, Pequignot F, Sempé M. Influence of body fat distribution during childhood on body fat distribution in adulthood: a two-decade follow-up study. *Int J Obes* 1990;14:473-81.
  33. Ross WD, Ward R. The O-Scale system. Rosscraft. Physique management systems Inc, 1984.